

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

2. REPORT TYPE  
Technical Papers

3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

5d. PROJECT NUMBER

2303

5e. TASK NUMBER

m2c8

5f. WORK UNIT NUMBER

345709

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)  
AFRL/PRS  
5 Pollux Drive  
Edwards AFB CA 93524-7048

8. PERFORMING ORGANIZATION  
REPORT

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)  
AFRL/PRS  
5 Pollux Drive  
Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S  
ACRONYM(S)

11. SPONSOR/MONITOR'S  
NUMBER(S)

please see attached

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

20030129 215

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

a. REPORT

Unclassified

b. ABSTRACT

Unclassified

c. THIS PAGE

Unclassified

17. LIMITATION  
OF ABSTRACT

A

18. NUMBER  
OF PAGES

19a. NAME OF RESPONSIBLE  
PERSON

Leilani Richardson

19b. TELEPHONE NUMBER

(include area code)  
(661) 275-5015

MEMORANDUM FOR PRS (In-House Publication)

FROM: PROI (STINFO)

11 May 2001

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-VG-2001-115**  
Mario Fajardo and Michelle DeRose, "Status of Cryosolid Propellants Task"

**AFOSR Molecular Dynamics Contractors' Meeting**  
**(Irvine, CA, 21 May 01) (Deadline: 18 May 01)**

**(Statement A)**

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review.

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b) appropriateness of references, if applicable; and c.) format and completion of meeting clearance form if required

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

4. This request has been reviewed by PR for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: \_\_\_\_\_  
\_\_\_\_\_

APPROVED/APPROVED AS AMENDED/DISAPPROVED

\_\_\_\_\_  
PHILIP A. KESSEL Date  
Technical Advisor  
Space and Missile Propulsion Division

# Status of CryoSolid Propellants Task

Mario E. Fajardo and Michelle E. DeRose

- \* Cryosolid Propellants Team
- \* HEDM Cryosolid Propellants Concept (Atoms in Solid Hydrogen)
- \* Cryosolid Propellants Payoffs, Objectives, Approach
- \* Requirement for Spectroscopic Diagnostics
- \* Rapid Vapor Deposition of Thick Parahydrogen (pH<sub>2</sub>) Solids
- \* Update on Al/pH<sub>2</sub> and Mg/pH<sub>2</sub> Experiments
- \* Opportunities for Supporting In-House Effort
- \* Recommendations for Future Experiments
- \* Open Discussion

# Cryosolid Propellants Team

- \* Mario E. Fajardo, Michelle E. DeRose, and Simon Tam  
Bill Larson and Jessica Harper (B atom source)
- \* J. Boatz, J. Mills, P. Langhoff, and J. Sheehy (in-house theory)
- \* FY00 Interactions with AFOSR Contractors:
  - P. Dagdigan @ Johns Hopkins: Al/H<sub>2</sub> & B/H<sub>2</sub> Complexes
  - M. Alexander @ U. Maryland: B/H<sub>2</sub> Interaction Potentials
  - G. Voth @ U. Utah: Path-Integral Monte Carlo Simulations
  - G. Scoles & K. Lehmann @ Princeton U.: Helium Clusters
- \* External Collaborators:
  - T. Momose @ Kyoto U.: High Resolution IR Spectroscopy
- \* Summer Visiting Professors:
  - R.J. Hinde @ U. Tennessee: Dopant-Induced IR Activity
  - D. Anderson @ U. Wyoming: Dopant IR Absorptions

# “Revolutionary” vs. “Evolutionary”

## HEDM Concepts

\* “Revolutionary” means better than LOX/LH<sub>2</sub>:

LOX/LH<sub>2</sub>       $\Delta H_{\text{sp}} = 12.6 \text{ MJ/kg}$  (3.0 kcal/g)

HEDM Target:       $\Delta H_{\text{sp}} > 15.0 \text{ MJ/kg}$  (3.6 kcal/g)

\* Early (c1990) Revolutionary HEDM Concepts:

tetrahydrogen (H<sub>4</sub>)

metastable triplet helium (He\* and He<sub>2</sub>\*)

spin-polarized atomic hydrogen (H↑)

high-spin species (<sup>5</sup>CO)

dications (AB<sup>++</sup>, ABC<sup>++</sup>)


♥ “non-metallics” (e.g. O<sub>4</sub>/H<sub>2</sub>, N<sub>4</sub>, N<sub>8</sub>, N<sub>20</sub>)  N<sub>5</sub><sup>+</sup>!

metallic hydrogen

♥ metal atoms and clusters in solid H<sub>2</sub>

# Cryosolid Propellants Concept

Use cryogenic solid hydrogen as a “packaging material” to store energetic species such as metal atoms and clusters.



# Cryosolid Propellants Payoffs

## Increased Specific Impulse

$$I_{sp} \propto \sqrt{\Delta H_{sp}}$$

$$\text{LOX/LH}_2 : I_{sp} = 400 \text{ s}$$

$$5\% \text{ B/sH}_2 + \text{LOX} : I_{sp} = 500 \text{ s (+25\%)*}$$

$$5\% \text{ Al/sH}_2 + \text{LOX} : I_{sp} = 450 \text{ s (+12\%)*}$$

\* calculated for  $P_{\text{chamber}} = 1000 \text{ PSIA}$ ,  $P_{\text{exhaust}} = 14.7 \text{ PSIA}$

## Greater Propellant Density

$$\text{liquid H}_2 @ 20 \text{ K} : \rho = 0.070 \text{ g/cm}^3$$

$$\text{solid H}_2 @ 2 \text{ K} : \rho = 0.087 \text{ g/cm}^3 (+25\%)$$

$$50/50 \text{ liquid He/solid H}_2 : \rho = 0.105 \text{ g/cm}^3 (+50\%)$$

# Atom Additive Payoffs (5 % molar)

Sea level specific impulse,  $I_{sp}$ , in seconds (% change)

$P_{\text{chamber}} = 1000 \text{ PSIA}$ ,  $P_{\text{exhaust}} = 14.7 \text{ PSIA}$

Additive	in standard state		as atoms		monoprop.	
	<u><math>M(5\%)/\text{LOX}/\text{H}_2</math></u>		<u><math>M(5\%)/\text{LOX}/\text{H}_2</math></u>		<u><math>M(5\%)/\text{H}_2</math></u>	
none	403					
C	381 (-5%)		515 (+28%)		515 (+28%)	
B	407 (+1%)		508 (+26%)		465 (+15%)	
Be	427 (+6%)		493 (+22%)			
Si	400 (-1%)		460 (+14%)			
Al	407 (+1%)		454 (+13%)			
H	403		430 (+7%)		380 (-6%)	
Li	404		428 (+6%)			
Mg	400 (-1%)		416 (+3%)			



# Cryosolid Propellants Objectives

- \* Make solid hydrogen samples (any size) containing 5% molar concentration of trapped energetic additives.
- \* Measure absolute concentrations of energetic species.
- \* Scale-up samples; produce  $\sim 1 \text{ cm}^3$  samples in our lab.

Example: 5% Al/pH<sub>2</sub>,  $V = 1 \text{ cm}^3$


assume each Al atom replaces one H<sub>2</sub> molecule

$\Rightarrow 58 \text{ mg Al} / 83 \text{ mg H}_2$  (\*see display item\*)


$\therefore \rho = 0.142 \text{ g/cm}^3$  (+100%)

# Cryosolid Propellants Approach

- \* Rapid vapor deposition of metal atom vapor and pre-cooled parahydrogen gas onto a liquid helium cooled substrate in vacuum.



# Recombination at High Concentrations



Bernoulli Trails (statistical) model of dopant agglomeration:  $P(k, 12) = \binom{12}{k} f^k (1-f)^{12-k}$

# Requirement for Spectroscopic Diagnostics

- \* Develop spectroscopic techniques to identify and measure concentrations of trapped species.
- \* Beer's Law:  $A = \sigma N d$   
 $A$  = absorbance =  $-\ln(I/I_0)$   
 $\sigma$  = absorption cross-section  
 $N$  = species number densities  
 $d$  = pathlength  
 $N d$  = "column density"
- \* UV/vis absorption for low metal atom column densities.
- \* IR absorptions, direct and dopant-induced  $pH_2$  transitions, for reaction products & metal atoms at large column densities.

# The Perils of Calorimetry

108

GEORGE C. PIMENTEL

TABLE IX  
CONCENTRATIONS OF FREE RADICALS REPORTED

Radical	Matrix	Mole per cent radicals	Method of production and estimate <sup>a</sup>	Reference
O	O <sub>2</sub>	4-20 <3 ~1	Gas, cal Gas, IR Gas, cal	Minkoff <i>et al.</i> (1959). Harvey and Bass (1958) Broida and Lutes (1955)
OH	Ca(OH) <sub>2</sub>	0.6	γ, ESR	R. Livingston <sup>b</sup>
N	N <sub>2</sub>	4	Gas, cal	Minkoff <i>et al.</i> (1959)
		0.2	Gas, cal	Broida and Lutes (1956)
		0.03	γ, ESR	Wall <i>et al.</i> (1959b)
		>0.03	Gas, cal	Fontana <sup>c</sup> (1958)
		0.01-0.04	Gas, MS	Matheson and Smaller (1955)
OH(?)	HCOOH	0.2	γ, ESR	Wall <i>et al.</i> (1959a)
CH <sub>3</sub>	CH <sub>4</sub>	0.14	γ, ESR	Wall <i>et al.</i> (1959a)
H	CH <sub>4</sub>	0.1	γ, ESR	Cole and Harding (1958)
N	NH <sub>3</sub>	0.1	Gas, ESR	Livingston <i>et al.</i> (1955)
H	HClO <sub>4</sub> -H <sub>2</sub> O	0.1	γ, ESR	Matheson and Smaller (1955)
H, NH <sub>2</sub> (?)	H <sub>2</sub> O	0.01	γ, ESR	Matheson and Smaller (1955)
ROH	NH <sub>3</sub>	0.01	γ, ESR	D. Ingram <sup>b</sup>
H	Alcohols	~0.01	UV, ESR	Wall <i>et al.</i> (1959a)
	H <sub>2</sub>	0.0006	γ, ESR	

<sup>a</sup> Abbreviations: gas = rapid condensation of gaseous radicals; γ = gamma ray *in situ* production; UV = photolytic *in situ* production; IR = infrared analysis; cal = calorimetry; MS = magnetic susceptibility.


<sup>b</sup> Private communication.

<sup>c</sup> Fontana, B. J. (1959). *J. Chem. Phys.* **31**, 148.

\* Difficult to distinguish energy release by small concentrations of very energetic species vs. low energy re-arrangements of host.

[A.M. Bass and H.P. Broida, "Formation and Trapping of Free Radicals" (Academic, New York, 1960)]

# Transmission Spectrum of Li/nH<sub>2</sub>, d $\approx$ 10 $\mu$



M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

# Optical Scattering in Solid Hydrogen

## Crystal Growing and Quality (p. 81)

“There is a considerable art to growing hydrogen crystals of high quality. Good crystals are always grown slowly from the melt; a rapid freeze from the gas produces snow.”


## Crystallite Light Scattering (p. 83)

“The reason that a good hydrogen crystal is so hard to see is its low refractive index...an estimated 1.16!

Yet a 1 mm-thick layer of hydrogen crystallites can be a completely opaque brown-black.”


[P.C. Souers, Hydrogen Properties for Fusion Energy (UC Press, Berkeley, 1986)]

# ortho- and para-hydrogen






# Rapid Vapor Deposition of Gram-Scale Optically Transparent pH<sub>2</sub> Solids (c1997)




M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).  
S. Tam and M.E. Fajardo, Rev. Sci. Instrum. **70**, 1926 (1999).

# Electronic Spectroscopy of B/pH<sub>2</sub> (d≈2 mm)




S. Tam, M. MacIer, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys. **113**, 9067 (2000).  
 [J.R. Krumrine, S. Jang, G.A. Voth, and M.H. Alexander, J. Chem. Phys. **113**, 9079 (2000)]

# Solid $\text{pH}_2$ Thickness from IR Spectra



S. Tam and M.E. Fajardo, Rev. Sci. Instrum., submitted (2001).

# Absorption Intensity vs. Thickness



new values:

$$\int A_{Q+S} d\tilde{V} / d_{\text{total}} = 82 \pm 2 \text{ cm}^{-2}$$

$$\int A_{S+S} d\tilde{V} / d_{\text{total}} = 6.3 \pm 0.3 \text{ cm}^{-2}$$

$$\tilde{\alpha}_{Q+S} = 4.84(\pm 0.13) \times 10^{-14} \text{ cm}^3/\text{s}$$


$$\tilde{\alpha}_{S+S} = 0.35(\pm 0.02) \times 10^{-14} \text{ cm}^3/\text{s}$$

literature (c1960):

$$\tilde{\alpha}_{Q+S} = 4.5(\pm?) \times 10^{-14} \text{ cm}^3/\text{s}$$

S. Tam and M.E. Fajardo, Rev. Sci. Instrum., submitted (2001).

# Constant pH<sub>2</sub> Deposition Efficiency



# High Flux HEDM Sources

- \* Purchased commercial Al evaporator; PBN crucible holds  $\approx 10$  g Al in horizontal orientation.
- \*  $T_{\max} = 1200\text{ }^{\circ}\text{C} \Rightarrow P_{\text{vap}}(\text{Al}) \approx 8 \times 10^{-3} \text{ torr} \Rightarrow \Phi_{\text{Al}} \approx 10^{18} \text{ \#/cm}^2\text{-s}$






Figure 1-3: Schematic of the EPI SUMO™ Effusion Cell.

# 60 ppm Al/pH<sub>2</sub> UV Absorption (d=0.14 mm)




# Al-induced IR Absorption Spectrum




Analysis in collaboration with Prof. R.J. Hinde, U. Tennessee (Knoxville).




# Correlation Between UV & IR Absorptions



# Constant Al Atom Deposition Efficiency




# UV Spectrum of Recombined Al/pH<sub>2</sub>



# Recombination/reaction in $\text{Al/pH}_2$




# Summary of Al/pH<sub>2</sub> Deposition Results



# 1000 ppm Al/Ar

\* Evaporate Al from 1 mm dia. Ta filaments



Al atoms in solid Ar

$$N_i = \frac{2.303 \ m_e \ c^2 \ \int A_{10} \ d\lambda}{\pi \ e^2 \ \lambda_0^2 \ f_{ik} \ d}$$

$$\int A_{10} d\lambda = 5.4 \text{ nm}$$

$$d = 4 \text{ } \mu\text{m}$$

$$N_{\text{Al}} = 2.8 \times 10^{19} \text{ } \#/\text{cm}^3$$

$$C_{\text{Al}} = 0.1 \%$$

$$\Delta t = 13 \text{ min}$$

$$\Phi_{\text{Al}} = 1.5 \times 10^{13} \text{ } \#/\text{cm}^2\text{-s}$$

$$(\text{a}) \ R = 5 \text{ cm}$$

## Al/pH<sub>2</sub> Summary

- \* Demonstrated trapping of  $\approx 0.1$  % Al atoms in thin Ar solids using home-made Al atom source.
- \* Demonstrated Al-induced IR absorption by pH<sub>2</sub> molecules as a diagnostic of Al atom concentrations in high column density samples.
- \* Attempts (four to date) to exceed 0.02 % Al atom concentration in thick pH<sub>2</sub> solids failed, resulting in catastrophic recombination (and reaction?) of the Al atoms.
- \* Interpretation: thick vapor deposited pH<sub>2</sub> solids impede dissipation of heat released upon atomic recombination, causing a recombination cascade. Attempt depositions of thin ( $\sim 10$   $\mu\text{m}$ ), high Al atom concentration Al/pH<sub>2</sub> samples (worked for Ar matrices).

# Is Catastrophic Recombination Ubiquitous to Rapid Vapor Deposited $pH_2$ Solids?

$$* \quad \frac{1}{k_{\text{recomb}}} = \frac{1}{k_{\text{intrinsic}}} + \frac{1}{k_{\text{diff}}}$$

\*  $k_{\text{intrinsic}}$  depends on M-M interaction potential,  
increases with increasing well depth and range

\*  $\Rightarrow$  try Mg atoms as “HEDM” dopant  
Mg-Mg and Mg-Mg<sub>n</sub> “van der Waals” interactions




# M-M Diatomic Ground State Potentials


\*\*\*\*\*INCOMPLETE SLIDE\*\*\*\*\*

- \* Will make PLOT comparing Al-Al and Mg-Mg ground state potentials, available from open literature.
- \* if not too cluttered, include Li-Li, B-B potentials, too.
- \* shallower wells and especially weaker long-range interactions mean reduced intrinsic recombination rates, according to G. Voth.

# Annealing of 10 ppm Mg/pH<sub>2</sub> (d=0.07 mm)



# As-Deposited 35 ppm Mg/pH<sub>2</sub> (d=0.02 mm)



# Contractor Support of In-House Effort

- \* Assignment of Al/pH<sub>2</sub> UV absorptions.  
can change calculated Al atom concentrations by factor of 2!
- \* Modeling of Al atom induced IR absorptions
- \* Modeling of Mg recombination in solid pH<sub>2</sub> (?)
- \* Open Discussion:  
coordination of efforts  
other suggestions?

# Recommendations for Future Experiments

## \* In-House:

Complete Al/pH<sub>2</sub> and Mg/pH<sub>2</sub> experiments

Compare Al vs. Mg to test effects of  $k_{\text{intrinsic}}$

Vary pH<sub>2</sub> deposition rate/sample thickness, and compare with

Ne and Ar hosts to test  $k_{\text{diff}}$

Evaluate feasibility of rapid vapor deposition method

Thermal B atom source SBIR project underway (B. Larson)

## \* AFOSR Contractors:

Cluster Pickup and Deposition

Deposition Directly onto lHe II (ala Gordon)

In-Situ Photolysis of HEDM Precursors (Apkarian, Stwalley)


Open Discussion: other suggestions?

# Future Direction: Cluster Deposition (?)

## Toward the Production of Measurable Quantities of Highly Doped Solid Hydrogen

High Energy Density Matter Contractors Conference  
June 4-7, 1995


Berton Callicott, Kenneth C Janda, V. A. Apkarian, R. B. Gerber  
Zhiming Li, P. Taborek and J. Rutledge  
Advanced Cryogenic Materials Group  
University of California  
Irvine, California



The goal of this effort is to produce measurable quantities of highly doped (5-10%) solid hydrogen. The desired dopant atoms are C, B, O, and N. Our approach to this material is via the cluster deposition technique, as illustrated in Figure 1. First, a beam of large He clusters (between  $10^4$  and  $10^5$  He atoms per cluster) is created following the methods of Toennies<sup>1</sup> and Scoles.<sup>2</sup> Next, the He clusters are passed through a scattering chamber that contains hydrogen molecules and, in the process, "pick-up" between 12 and 20 hydrogen molecules. During the pick up process, sufficient He atoms evaporate for the cluster to maintain a temperature of about 1 K, and the hydrogen is expected to make a small micro-crystal within the He cluster. Next, the He cluster picks up the desired dopant atom. Again, He atoms evaporate to cool the cluster back to 1 K.

- \* Most excess heat is dissipated before the clusters are deposited.
- \* Chance to beat the "statistical limit" of stored atom concentration.
- \* Either higher fluxes or UHV deposition environment required.

# Future Direction: Deposition onto IHe II (?)



Based on E.B. Gordon's work

“Big-Flush” (c1995)

CESE discharge of N<sub>2</sub>/He

Optical emission was only diagnostic  
(N/D<sub>2</sub> samples didn't glow!)

Need alternative (species specific)  
experimental diagnostics: ESR,  
NMR, FIR absorption... (?)


# Future Direction: In-Situ Photolysis (?)

\* V.A. Apkarian, et al. experiments (c1994):



(5% O/H<sub>2</sub>  $\Rightarrow$  25s (6%) I<sub>sp</sub> improvement, so Al > O > H)  
 UV induced desorption of H<sub>2</sub>O inside vacuum chamber  
 complicates monitoring O + H<sub>2</sub> reactions.

\* W.C. Stwalley proposal (c1992):





# Experimental Admonitions

- \* Must work with hydrogen! Some efforts on model systems are fine, but results may not generalize cleanly to hydrogen.
- \* Must focus on production and quantitative measurement of  $\sim 1\%$  concentrations of energetic species in solid hydrogen. Species specific diagnostics are preferred.
- \* Worry about scaling up later, must demonstrate progress towards larger concentrations to maintain viability of Cryosolid Propellants Task.